

Contents lists available at ScienceDirect

Heliyon

journal homepage: www.cell.com/heliyon



Research article

Does agricultural ecology cause environmental degradation? Empirical evidence from Bangladesh



Shanjida Chowdhury ^a, Sunjida Khan ^b, Md Fouad Hossain Sarker ^{c,*}, Md Kabirul Islam ^d, Maruf Ahmed Tamal ^e, Niaz Ahmed Khan ^f

- ^a Department of General Educational Development, Daffodil International University, Dhaka, Bangladesh
- ^b Department of Business Administration, Daffodil International University, Dhaka, Bangladesh
- ^c Department of Development Studies, Daffodil International University, Dhaka, Bangladesh
- d Department of Multimedia and Creative Technology, Daffodil International University Dhaka, Dhaka, Bangladesh
- ^e Division of Research, Daffodil International University, Dhaka, Bangladesh
- f Department of Development Studies, University of Dhaka, Dhaka, Bangladesh

HIGHLIGHTS

- The relationship between agro-economic climate and CO₂ emissions in Bangladesh from 1985 to 2017 is analysed
- The results exhibit that production of cereals, other agricultural products, and land production, increased CO₂ emissions.
- Unplanned, static crop intensity and a lack of modern farming infrastructure degrade the environment.
- The findings may help Bangladesh achieve UN-SDG targets 7, 9, 12, and 13.

ARTICLE INFO

Keywords: Environmental degradation Agricultural ecology CO₂ emission Bangladesh

ABSTRACT

Agricultural sector accelerates a nation's economic growth towards sustainable development. There exists a significant relationship between agriculture and the environment. Sustainable agricultural development ensures food quality and in tandem prevents natural calamities like drought. However, in order to fulfill the food demand of a growing population, poor law quality and untenable agriculture practices arise, which in turn lead to environmental degradation. The current study explores the relationship between the agro-economic atmosphere and CO2 emissions as a measure of environmental degradation in Bangladesh between the years of 1985 and 2017. To exhibit the long-run relationship of agricultural ecology and carbon dioxide emissions, three cointegrated equations- Fully-modified ordinary least square (FMOLS), Dynamic ordinary least square (DOLS), and Canonical cointegrated regression (CCR) were assessed. For cointegration, Bayer-Hanck cointegration was implied. In long-run estimates, it was found that livestock, rice area harvested, cereal production, and other crop production impeded environmental dilapidation. The Granger Causality Test enabled unidirectional causality towards burned biomass (crop residues), the agricultural economy, and carbon emissions. Therefore, this dimension's causality concluded that carbon dioxide emissions were caused by cereal production, other agricultural production, and agricultural land production. The overall findings of this study could potentially assist the Government of Bangladesh and the necessary authorities for implementing synchronized policies to help reduce environmental pollution and set an example for other developing nations like Bangladesh.

1. Introduction

Environmental degradation arises mainly due to the continuous emission of greenhouse gases from anthropogenic activities [1]. The Rio

(1992) Earth Summit inferred that no credible alternative exists apart from emphasizing social, economic, and environmental dimensions to save the earth, and these three factors were closely interdependent with each other [2]. In order to meet a burgeoning population, modern

E-mail address: fouadsarker@daffodilvarsity.edu.bd (M.F.H. Sarker).

^{*} Corresponding author.

technologies manipulate the ecological atmosphere, thereby leading to environmental degradation. Advanced technologies such as biomass have a negative impact on the environment [3]. Additionally, using modern technologies leads to consequences such as deforestation, land degradation, soil erosion, water pollution, air pollution, etc. which in turn affect farming and farm produce [4]. Since population is one of the most important determinants of demand, a continuous increase in the world population increases the worldwide demand for food to meet basic human needs [5].

Agriculture is a leading form of global land management and the agricultural ecosystem covers nearly 40% of the land surface on earth [6]. Agricultural ecosystems are significantly limited to rural areas, with almost 50% of the world's population living in these rural areas. Bangladesh is no different with approximately 70% of the population living in villages and depend on agriculture for their livelihood [7]. In many developing countries, agriculture is considered as the primary driver of economic growth, with almost 75% of agricultural value addition being produced in these nations [8]. Agro-ecosystems are the natural processes or ecosystems on agricultural land which have been altered to sustain the production of agricultural products such as food, harvestable goods, and fiber [9]. It is designed by humans and is used for enhancing agricultural production. Rapid food and fiber production are also considered as the causal factor for environmental degradation [10] (see Figure 1).

To ensure environmental quality for survival, it has been emphasized to maintain the same through sustainable and appropriate use of available resources. Land, water, mining, soil, forests, etc. are considered some common natural resources on which all agricultural activities rely on. Bangladesh has a rapidly expanding population and must produce agricultural products quickly and import from other countries to meet their needs [11]. Technology, as well as the judicious utilization of chemicals, biomass, and other materials would aid in the development of speedy results. Agricultural operations such as land reclamation, irrigation, crop breeding, and deforestation among other things end up polluting the environment [12]. Rapid farming activities may cause soil degradation, soil erosion, air pollution, deforestation, desertification, and many other negative effects, which in turn would affect farming and farm produce [12]. Increased pesticide usage harms farmworkers via extended exposure times and contaminate ground and surface water sources, including harming downstream users and inland fisheries [13]. According to a recent World Bank report, Bangladesh is one of the most densely populated countries in the world [14]. This has causes a significant rise in the demand for diverse agricultural products which has resulted in continuous greenhouse gas (GHG) emissions. According to a report by Climate Watch [15], GHG emissions in Bangladesh increased significantly and reached a historic high of 220.75MT CO2e in 2018. The increase in emissions was mainly driven by agriculture (88.53MT) followed by Energy (85.84MT), Land-Use Change and Forestry (21.78MT), Waste (20.64MT) and Industrial Processes (3.97MT). Though meeting the requirements of a rising population is vital, a healthy environment is also a significant requirement for human survival [16]. Based on the review of past research, it was concluded that more insight into the relationship between agricultural ecology and environmental degradation in Bangladesh was needed. Numerous studies have been conducted in Bangladesh focusing on environmental degradation, but the relationship between environmental degradation and agricultural ecosystems has received little to no attention.

Based on the aforementioned points, the major objective of this study was to investigate the nexus between Bangladesh's agro-economic environment and $\rm CO_2$ emissions as a measure of environmental degradation from 1985 to 2017.

2. Literature review

Climate change is the most pressing issue facing the entire globe today. Anthropogenic greenhouse gas emissions contribute to global warming, consequently causing long-term impact on weather patterns and ozone layer depletion. Agriculture services such as land erosion control, irrigation water management, insect infestation control, boosting nutrient cycling, pollination, and others are utilized to increase harvestable product yield. However, these processes interfere with the natural cycle and are therefore considered detrimental due to causing water loss and soil quality reduction amongst other harmful consequences [17].

Agriculture operations have a net negative impact on overall environmental quality [18]. The waste produced during these activities pollute the soil, water, and environment. A past study discovered a positive association between agricultural productivity and ecological pressure, i.e. as the product grows, so does the demand on the environment. Following the Green Revolution, synthetic fertilizers and pesticides have accounted for a substantial percentage of agriculture dependency [19]. Rapid food production with synthetic fertilizers is preferred, not only because of the importance of agriculture but also to alleviate food safety concerns. Consequently, it has caused hazards for human health and the surrounding survival environment. Another past study [20] identified a U-shaped relationship between carbon dioxide emission (CO₂) and agricultural development. The analysis was based on the agriculture sector in Ghana and recommended some policy suggestions emphasizing large-scale adoption of environment-friendly techniques for reduction of emissions and concurrent agricultural development. It has also been noted [21] that wetland rice production acted as an important contributor in reducing greenhouse gas (GHG) emissions. The concerned study indicated that non-puddling rice production with increased residue retention reduced more GHG emissions in comparison to non-puddling rice production with decreased residue retention. Therefore, high residue retention contributed to increased emissions removal. Another study revealed a positive relationship between carbon dioxide emissions (CO₂) and agricultural development, leading to policy recommendations that focused on large-scale adoption and ecologically acceptable approaches for decreasing emissions and advancing agricultural development [20]. Past studies have shown that puddling with lower residue retention during rice production reduces GHG emissions more when compared to non-puddling methods [21]. Consequently, it can be stated




Figure 1. Benefits and concerns of diversified agroecosystems [Source: FAO AQUASTAT (2009)].

that high residue retention has a greater influence on decreasing emissions.

Rahman [22] performed a study to establish an environmentally beneficial technique known as 'Environment Smart Agriculture' (ESA). The study used an indicator to exhibit that using chemicals had a positive impact on agricultural produce's long-term viability and also stated that there had to be an improvement in the farmers' understanding of the use of chemicals in agriculture. Another study [23] devised an indicator for determining the level of environmental degradation. The results of the study were indicative that the production of HYV rice caused a 27-69% increase in the theoretical maximum amount of environmental harm, alluding to substantial environmental deterioration and thereby raising worries among policymakers. According to a study by Faroque [24], the indiscriminate usage of chemicals in agricultural production caused environmental deterioration, which in turn posed a threat to agricultural viability. The government has been attempting to promote on-farm resources and encourage alternative usage of external inputs and pesticides. The study stressed the importance of reducing chemical use in agriculture.

Higher economic growth is dependent on the extensive use of resources to sustain demand, and this can create serious environmental pollution. Thus, it can be stated that there is a trade-off relationship between environmental pollution and economic growth. Underdeveloped and developing nations mostly depend on agriculture for boosting their economy. As and when their agricultural productivity increases, more resources tend to be used, thereby leading to environmental degradation [25]. It has been exhibited that the consumption of natural resources via mining, agriculture, and deforestation has the potential to degrade environmental quality by increasing emissions [26]. The ambition to expand the agricultural frontier in Livelihood Adaptation to Climate Change (LACCs) where agricultural activities are dominating leads to damaged vegetation, desertification, and deforestation, exposing the land to deterioration [27].

Nathaniel and Adeleye [28] showed that the UN-SGDs (SDGs-9) emphasized the need for reducing carbon emissions and preventing environmental damage. Carbon emissions significantly contribute to environmental degradation, thereby posing a challenge to establishing and maintaining a sustainable ecosystem. The authors also studied a variety of factors that influenced CO₂ emissions and their potentially harmful consequences on the ecosystem. Usman et al. [29] discovered a substantial link between agricultural growth, economic growth, and the use of non-renewable energy, which caused environmental degradation. The study also found a link between carbon emissions and economic growth, which was a roadblock to meeting sustainable development targets.

According to Hafeez et al. [30], agriculture was responsible for 21% of global CO_2 emissions, and concluded that agriculture and energy consumption were driving environmental degradation. They also emphasized energy efficiency and use of eco-friendly technology. Olanipekun [31], stated that growing population and poverty had a significant impact on environmental degradation in emerging countries. As the population grows, so does the demand for food, necessitating massive food production in a short period of time. Consequently, agricultural practices relying on rapid output may be unsustainable and negatively impact the environment due to CO_2 emissions.

The natural climate system's balance is usually stabilized by the local ecosystem, including carbon, nitrogen, and water cycles. Greenhouse gases have been shown to be the long-term sources of climate change [32]. Global warming results from increased greenhouse gas emissions and it has the potential to alter the environment over time [33]. CO₂, F, CH₄, and N₂O are the main greenhouse gases found in anthropogenic emissions and among them, CO₂ contributes to nearly 76%, which causes global warming [34]. Environmental degradation has been of concern worldwide over the previous decades [35], whereas and the agricultural sector accounts for 10–15% of global emissions [36]. Even though developing countries heavily rely on agriculture, rising CO₂ emission levels have been shown to reduce land productivity and inversely affect agricultural output [37].

CO₂ emissions have adversely changed the climate, thereby disrupting the agro-ecological balance and income distribution, which creates insecurity in food consumption [38]. Agricultural greenhouse gases are mainly created from the cultivation of rice, crop residue burning, and livestock. Large volumes of GHGs are emitted during agricultural production mainly due to overexploitation of agricultural resources such as land, usage of chemical fertilizer and pesticides, ultimately resulting in climate change. Additionally, studies have also noted that rising food production to feed the world's growing population has resulted in inefficient farming methods that inevitably degrade the natural environment [39]. Agriculture has also been shown to be significantly impacted by environmental challenges such as climate change, global warming, agricultural land degradation, water and air pollution, and biodiversity loss. Thus, future expansion of food output must be balanced against the growing scarcity of natural resources [36]. Studies have also shown that CO₂ emissions from agricultural sources account for approximately 21% of total anthropogenic GHG emissions [38]. As a result of changing agro-ecological conditions, environmental degradation ultimately reduces agricultural output, leading to unequal income distribution which in turn leads to food insecurity [38].

Sustainable Development Goals (SDGs) formulated by the United Nations is a global policy action for all nations to implement in order to promote prosperity while simultaneously protecting the environment. Accordingly, ensuring a poverty-free world by the year 2030 forms the first goal of the SDGs [36]. The SDGs aim to increase agriculture productivity and ensure stable food security for all people by promoting sustainable consumption, production, and distribution approaches. Consequently, agriculture forms a vital issue for improving the quality of life and achieving the SDGs [40]. It is considered to be possible to reduce the inverse effect of agriculture on the environment by adopting strong and appropriate policies and technological transmission [41, 42].

According to Roberts [43], human communities have created a threat and negatively changed the global ecosystem via increased CO_2 emissions. The increase in the population and animals of developing nations increases the production of food and livestock and thereby leads to increased emissions [44]. To meet the global population's food demand, the agriculture sector has been under enormous strain and has produced emissions [45]. Sustainable agriculture helps increase food production while concurrently reducing the use of pesticides. Agriculture production has also been observed to accumulate carbon at times, thus polluting the environment [46]. It has been shown that there is a significant relationship between economic growth and environmental pollution. On the other hand, pollution reduction and environmental balance are not prioritized enough in developing and underdeveloped nations [47, 48].

2.1. Agriculture and environment in Bangladesh context

Bangladesh is primarily an agricultural country and the agricultural industry plays a vital role in driving the economic growth of the country. Agriculture employs over half of the Bangladeshi population, with more than 70% of the country's land dedicated to agricultural production [49]. According to the report [50], agriculture's contribution to the GDP of Bangladesh grew to 11540.50 million BDT in 2021 from 11242.30 million BDT in 2020. However, the agricultural sectors (e.g., rice cultivation, enteric fermentation, synthetic fertilizers, livestock manure etc.) contributed to about 28% of total GHG emissions in 2018 [51], thereby posing a huge environmental concern. Similar findings were obtained in an earlier study [52], where the authors found a significant relationship between agricultural development and CO₂ emissions. The research also showed that population, agriculture energy intensity and the economic activities were the factors most responsible for the increase in CO2 emissions which negatively impacted agriculture as well as the environment. In another study [53], the environmental consequences of Green Revolution technology diffusion in Bangladesh was explored. The study used 21 villages in three agro-ecological zones in Bangladesh as the sample area and factors like loss of soil fertility, trends in fertilizer and

pesticide productivity at the national level, as well as farmers' awareness etc. were selected and examined as evidences. The results of the study indicated that Bangladesh had lost soil fertility in 11 out of its 30 agro-ecological zones to the tune of 10–70% between 1968 and 1998 due to intensive crop cultivation practices. A later study [54] revealed a casual nexus between agricultural output and CO_2 emission levels wherein empirical evidences revealed that the agricultural output of Bangladesh was not a Granger causal for CO_2 emissions, but the country's CO_2 emissions were a Granger causal for its agricultural output. Similarly, in another study [55], the authors identified that enormous population, economic factors, and human activities of low-income level nations like Bangladesh generated increased levels of CO_2 emissions. Currently, Bangladesh contributes to around 0.14% of the world's CO_2 emissions.

Using the cointegration technique, this study proposes to investigate the causal relationship between agricultural ecosystems and environmental degradation. To the best of the authors' knowledge, none of the past Bangladesh centric studies have examined the causal relationship between agricultural ecosystems and environmental contamination. This study envisages to be a major step forward in environmental and agricultural research.

3. Materials and methods

This study investigated the linear relationship between environmental degradation and agricultural ecology in the period of 1985–2017 in Bangladesh by using annual datasets. Data, except for Bangladesh's carbon dioxide emissions, was collected by the FAO [56]. Due to the lack of availability of most recent data, this study was limited to data available until 2017. Table 1 highlights the description alongwith a variable name. Various studies have shown that biomass-burned crop residues [57], agriculture value-added [58], CO₂ equivalent to N₂O emissions from synthetic fertilizers [59], livestock stock [60], agriculture machinery [60], cereal production [60], all have a significant impact on greenhouse gas emissions. Based on these past studies, all of these variables were used as explanatory variables in this study.

The Intergovernmental Panel on Climate Change (IPCC) adopted the Tier 1 approach for measuring CO₂ emissions due to climate change. All member countries estimate their CO2 emissions under the guidance of this strategy [61]. The current study calculates this in kilotons (Kt) and as a dependent variable, and the remaining variables treated as independent variables. The value added to agriculture was calculated in terms of its percentage share of the GDP. Crop residues, cereal production, and other crop production were all taken into account and their units are represented in tons. Synthetic fertilizer emissions were quantified as gigagrams of CO2 equivalent (N2O). In the case of livestock capacity, the head number was counted for this study. Machinery and tractors used for agricultural purposes were calculated in raw numbers. Since rice is the core factor of agricultural production in Bangladesh, paddy harvested per hectare was considered for this study's purposes. In order to avoid multicollinearity, all variable data was converted to natural logarithmic form and the following regression was provided to exhibit the relationship between carbon emission and the explanatory variables.

$$\mathit{CO}_2 = \! f(\mathit{BM},\ \mathit{AG},\ \mathit{SF},\ \mathit{LS},\ \mathit{MC},\ \mathit{RI},\ \mathit{CP},\ \mathit{OT})$$

For exhibiting this linear combination, the equation was converted into the log-linear model. The final regression model for employing multiple cointegrated relationships is represented as follows:

$$\begin{split} &\ln(CO_2)_t = \beta_0 + \beta_1 \ln\!BM_t + \beta_2 \ln\!AG_t + \beta_3 \ln\!SF_t + \beta_4 \ln\!LS_t + \beta_5 \ln\!MC_t \\ &+ \beta_6 \ln\!RI_t + \beta_7 \ln\!CP_t + \beta_8 \ln\!OT_t + \varepsilon_t \end{split}$$

In this final equation, β_0 is the constant, β_1 to β_8 are the parameter coefficients of respected variables and ε_t is the stochastic error term. To anticipate the direction or effect of variables on carbon emission, it is dependent on the sign of β_i 's. for more granularity, biomass residues

Table 1. Definition of variables under econometric analysis.

Variable	Description
CO_2	Log of carbon dioxide emissions (Kt)
ВМ	Log of biomass-burned crop residues (dry matter; tons)
AG	Log of agriculture value added (% of GDP)
SF	Log of emissions of CO ₂ equivalent of N2O from synthetic fertilizers (Gg)
LS	Log of stock of livestock (number of head)
MC	Log of agricultural machinery (number of tractors)
RI	Log of rice area paddy harvested (Ha)
CP	Log of cereal production (tons)
OT	Log of other crop productions (tons)
LS MC RI CP	Log of stock of livestock (number of head) Log of agricultural machinery (number of tractors) Log of rice area paddy harvested (Ha) Log of cereal production (tons)

accelerate environmental degradation if $\beta_1>0$ and vice versa if <0. The agricultural output would increase if $\beta_2>0$ through the positive impact of carbon emission. Similarly, synthetic fertilizer and the amount of livestock would increase if β_3 and β_4 are positive. For exhibiting the proxy of modern technology on agricultural production, this study takes into account agricultural machinery as an explanatory variable. If the impact of the same is positive, then $\beta_5>0$, otherwise it decreases while being negatively impacted. To check the robustness of agricultural output, three more variables on crops-related production were employed for this study viz. production of principal food rice, cereal or crop and miscellaneous crops production. The linkage between environmental disaster and these three variables would be positive if β_6 , β_7 , and β_8 are individually positive.

We used FMOLS, DOLS, and CCR estimators to demonstrate long-term behavior. These three estimators are emphasized for the following reasons. Firstly, though the OLS estimator is super-consistent [62, 63], the estimated t-statistic becomes approximately normal at normal level or stationary. Moreover, the bias convergence turns low in finite samples. Secondly, the t-statistic becomes insignificant in the case of omitted dynamics captured by residuals and it suffers from serial correlation and heteroscedasticity. Thirdly, FMOLS and DOLS have specific criteria. FMOLS works on the I(1) series [64], rectifies endogeneity and serial correlation effects [65], while DOLS is employed to estimate long-run equilibrium which is corrected for potential simultaneity bias among regressors [65]. Finally, DOLS is more useful to Johansen's cointegration, as compared to FMOLS, or CCR. Being a robust but single equation method, misspecification error does not arise [62]. Moreover, DOLS has no demanding restriction on the order of variables, given that regressors should be integrated in order (1). DOLS regresses one of the I(1) variables on other I(1) variables, I(0) variables, and lags and leads to the first differences of the I(1) variables.

Unit root tests are considered important for performing time series analysis when arranging variables in a stationary order. This study employed a version of the unit root test known as Augmented-Dickey Fuller test [66] and Phillips-Perron [67] test to observe the order of integration. This study also used the Ng and Perron test which is considered as an effective modification of unit root tests. The inclusion of the PP test in this analysis was mainly due to its robustness to a set of time-dependent serial correlations and heteroscedasticities. The Akaike information criteria was used to evaluate the lag length in the ADF test case, while the Newey-West Bartlett kernel was used to select the bandwidth for the PP test. A conventional integration approach was applied and the null hypothesis generated suggested that the series was non-stationary, while it was stationary in the case of alternative hypotheses. The intercept and deterministic trends of time series data integrations were also used in this study.

3.1. Cointegration

Over the years, many cointegration methods have evolved. Engle and Granger [68] recommended a cointegration evaluation based on

projected long-term regression model residuals. Several cointegration experiments were developed much later, such as Johansen's system-based test [69], Boswijk's ECM-based F-test [70], and Banerjee et al.'s ECM-based t-test [71]. These cointegration approaches have distinct theoretical bases and produce contradictory findings, thereby creating a limitation. The effectiveness of ranking cointegration techniques is significantly dependent on the value of nuisance estimators [72]. This study used a newly established cointegration test proposed by Bayer and Hanck [73] to verify the existence of cointegration relationship between environmental destruction and agricultural ecology in Bangladesh to solve the power of the cointegration tests. A distinctive feature of this test is that it helps integrate multiple individual findings from the test into a more decisive outcome. Bayer and Hanck [73] recommended combining the individual cointegration test's computed significance level (p-values) with the following formulae postulated by Fisher (1 and 2):

$$EG - J = -2[\ln(p_{EG}) + \ln(p_{J})] \tag{1}$$

$$EG - J - B - BDM = -2[\ln(p_{EG}) + \ln(p_J) + \ln(p_B) + \ln(p_{BDM})]$$
 (2)

Where p-values of several distinct cointegration tests such as Engle-Granger [69]; Boswijik [70] and, Banerjee et al. [71] are exhibited by P_{EG} , P_{J} , P_{B} and P_{BDM} respectively. Fisher statistics of the estimated values when exceeding the critical values proposed by Bayer and Hanck [73] for the null hypothesis of no cointegration can be rejected.

3.2. Long-run estimates

The Fully Modified Ordinary Least Squares (FMOLS) estimator developed by Phillips and Hansen [74] and the Dynamic Ordinary Least Square (DOLS) proposed by Stock and Watson [75] have several advantages over OLS estimators. FMOLS is mainly applied for a semi-parametric approach for long-run parameter estimation [76, 77]. FMOLS provides consistent estimator parameters [72, 78] and transforms data and parameters [72] in the case of small samples and succession over endogeneity or serial measurement error, omitting biased variables, solving heteroscedasticity, or serial correlation. FMOLS estimator Adom [79] is as follows (3):

$$\widehat{\beta}_{FMOLS} = \left[\sum_{t=1}^{T} (x_t - \overline{x_1})(x_t - \overline{x_1})\right]^{-1} * \left[\sum_{t=1}^{T} (x_t - \overline{x_1})y_t^* + T\tilde{\triangle}_{EM}\right]$$
(3)

$$\mathbf{y}_{t}^{*} = \mathbf{y}_{t} - \widehat{\Omega_{EM}} * \Omega_{E}^{-1} \Delta \mathbf{x}_{t}$$

$$\widetilde{\Delta}_{EM} = \Delta_{EM} \widehat{\Omega_{E}^{-1}} \Omega_{EM}$$

Here, y_t^* is the transformed variable of the dependent variable to attain the endogeneity equation, and $\tilde{\Delta}_{EM}$ is the serial correlation correction.

DOLS is an extension of the Stock and Watson's [76] estimator, which is obtained from the following equation (4) -

$$y_t = \alpha_i + \beta x_t + \sum_{i=1}^k c_i \Delta x_{t-i} + v_{it}$$
(4)

 c_j is the coefficient of a lead or lag and is the first differenced explanatory variable. The estimated coefficient of DOLS is given by the following equation (5):

$$\widehat{\beta}_{DOLS} = \left[\sum_{t=1}^{n} q_t \ q_t \right]^{-1} * \left[\sum_{t=1}^{n} q_t \ y_t^* \right]$$
(5)

Here, q is {2(q+1) \times 1} vector of regressors and. $q_t = x_t - \overline{x_i} \Delta x_{t-q}$, Δx_{t+q}

Canonical cointegrated regression (CCR), introduced by Park [80], works similar to FMOLS. However, there is a difference between these two models. CCR works on data transformation only, while FMOLS converts data and parameters [79, 80]. Further, CCR is also a single equation-based regression, which can also be applied to multivariate regression. The CCR estimator is obtained as follows (6) from Adom [79]:

$$\widehat{\beta}_{CCR} = \left[\sum_{t=1}^{T} (x_t - \overline{x_1})(x_t - \overline{x_1}) \right]^{-1} * \left[\sum_{t=1}^{T} (x_t - \overline{x_1}) y_t^* \right]$$
(6)

3.3. Causality test

Changes in one variable may precede or follow changes in another variable, resulting in a variety of causal relationships. The objective of studying this causality is referred to as Granger causality (1969). The following models were studied to clarify the technique for analyzing Granger causality:

$$Y_{t} = \sum_{i=1}^{m} a_{i} Y_{t-i} + \sum_{i=1}^{m} b_{i} X_{t-i} + u_{t}$$
 (7)

$$X_{t} = \sum_{i=1}^{m} c_{i} X_{t-i} + \sum_{i=1}^{m} d_{i} Y_{t-i} + \nu_{t}$$
(8)

Based on the above models (7 and 8), the following cases are distinguished:

- The coefficients b_i of the variables X_{t-i} in (1) are statistically significant, while the coefficients c_i of Y_{t-i} in (2) are statistically different from zero. In this case, there is Granger causality from X to Y.
- The coefficients b_i of the variables X_{t-i} in (1) are not statistically significant, while the coefficients c_i of in Y_{t-i} (2) are statistically different from zero. In this case, there is Granger causality from Y to X.
- The coefficients of b_i variables X_{t-i} in (1) and c_i coefficients of Y_{t-i} in (2) are statistically different from zero. In this case, there is bidirectional Granger causality.
- The coefficients of b_i variables X_{t-i} in (1) and c_i coefficients of Y_{t-i} in (2) are statistically different from zero. In this case, there is Granger independence.

Consequently, Granger causality estimation examines the null hypothesis that one variable does not cause another variable by examining the statistical significance of the estimated coefficients.

4. Results and discussion

The present study attempts to observe the causal relationship between agricultural production and carbon emissions by using time series data from the year 1985 till 2017. Results of basic statistical analysis along with their correlation analyses are tabulated in Table 2. It can be observed that the mean, median, and mode are close to each other following natural logarithmic transformation. Skewness describes the thickness or flatness, whereas kurtosis describes the peak nature of the distribution. From the results obtained in this study, it can be observed that CO₂ emissions, the CO₂ emission equivalent for N₂O from synthetic fertilizer (SF) usage, and diverse crop production (OT) were all to the left. In the kurtosis analysis, none was distributed approximately, except for the CO₂ emission equivalent that comes from synthetic fertilizer usage. Almost all the variables rejected normality under the Jarque-Bera statistic. Correlation describes the degree and direction of association amongst variables. Agricultural value-added contribution to the economy (AG) relates negatively to all variables except for other crop production (OT). Also, for CO2 emissions, this variable's relationship was found to be more potent with burning of crop residues and biomass, agriculture machinery, cereal production, area harvested by rice

Table 2. Basic statistics with correlation matrix of variables.

	CO_2	BM	AG	SF	LS	MC	RI	CP	OT
Mean	6.761	15.652	3.104	8.661	16.664	8.654	16.18	17.427	12.988
Median	7.134	15.64	3.117	8.768	16.637	8.632	16.213	17.453	13.018
Maximum	8.23	15.732	3.547	9.051	16.874	8.838	16.3	17.876	13.412
SD	4.595	15.573	2.642	7.772	16.425	8.497	16.108	16.994	12.481
Skewness	1.188	0.051	0.279	0.334	0.134	0.102	0.061	0.299	0.207
Kurtosis	-0.491	0.213	0.075	-1.128	0.038	0.3	0.305	0.001	-0.439
Jarque Bera	1.897	1.741	1.657	3.412	1.89	1.815	2.092	1.567	2.847
Probability	3	2.429	2.512	7.231	1.703	2.425	1.646	2.822	1.094
Sum	0.223	0.297	0.285	0.027	0.427	0.297	0.439	0.244	0.579
Sum Sq. Dev.	223.115	516.528	102.417	285.818	549.926	285.576	533.949	575.085	428.602
Obs.	45.167	0.085	2.496	3.579	0.574	0.336	0.118	2.859	1.374
Correlations Matrix									
	CO_2	BM	AG	SF	LS	MC	RI	CP	OT
CO_2	1	0.725	-0.844	0.751	0.824	0.859	0.636	0.85	-0.198
BM	0.725	1	-0.813	0.633	0.782	0.839	0.852	0.884	-0.362
AG	-0.844	-0.813	1	-0.882	-0.968	-0.973	-0.767	-0.973	0.117
SF	0.751	0.633	-0.882	1	0.888	0.843	0.625	0.871	0.196
LS	0.824	0.782	-0.968	0.888	1	0.987	0.713	0.961	-0.049
MC	0.859	0.839	-0.973	0.843	0.987	1	0.748	0.975	-0.132
RI	0.636	0.852	-0.767	0.625	0.713	0.748	1	0.813	-0.344
CP	0.85	0.884	-0.973	0.871	0.961	0.975	0.813	1	-0.166
ОТ	-0.198	-0.362	0.117	0.196	-0.049	-0.132	-0.344	-0.166	1

production, and the CO_2 emissions equivalent for N_2O from synthetic fertilizers usage. A high correlation was observed among rice-harvested areas, cereal production, livestock, and agricultural machinery. Multicollinearity was also not observed.

4.1. Unit root test

A unit root test determines whether a time series component is nonstationary [81]. The results of unit root test presented in Table 3 highlights the effects of the ADF and the PP unit root tests. In both cases, the study variables were observed to be stationary at the first difference after logarithmic transformation, which was concurred to the findings of Ghimire et al. [82].

4.2. Ng-Perron test

This study also employed the Ng-Perron test for further investigation rather than relying only on the PP test and ADF test for stationarity testing. The goal of using this test was to show that Ng-Perron test was more powerful and more suggestive than the ADF test in the case of small sample size since the ADF test has been shown to be unreliable for small

samples [83, 84]. The results of the Ng-Perron test (at level and at first difference) are presented in Tables 4a and 4b. From the results, it can be seen that only OT was weakly stationary at this level, but all were statistically significant at the first difference. As a result, it can be stated that at the first difference or integration, each variable is significant.

4.3. Cointegration analysis

Fisher's statistics were used in this study to observed if the underlying variables were cointegrated. The integrated cointegration measures, including the EG-JOH and EG-JOH-BO-BDM measures, are highlighted in Table 5. EG-JOH and EG-JOH-BO-BDM's F-values were found to be greater than a critical value, implying that the null hypothesis of no cointegration can be rejected. A similar result was obtained by Chaouachi and Balsalobre-Lorente [85] in an earlier study. Therefore, Johansen and Bayer-Hanck's cointegration tests confirmed that a long-run cointegration did occur between the selected variables in Bangladesh for the time period of 1985–2017.

Bayer and Hanck [73] developed a combined cointegration approach that strengthens the cointegration analysis. This innovative technique amalgamates findings of previous cointegration and depicts on Fisher

Table 3. Result of Augmented-Dickey Fuller (ADF) test and Phillips-Perron (PP) test.

Variable	ADF Test				PP Test				
	At level		At difference	At difference		At level		At difference	
	С	C + T	С	C + T	С	C + T	С	C + T	
CO_2	0.267	-1.975	-3.509***	-3.306***	-2.232	-3.215	-16.305***	-15.691***	
ВМ	-0.056	-3.202	-3.012**	-3.162***	-0.921	-2.813	-6.620***	-6.708***	
AG	0.062	-3.065	-5.549***	-3.969***	-0.051	-3\2.095	-6.763***	-6.643***	
SF	-2.749	-2.910	-4.554***	-5.525***	-5.570	-2.207	-6.393***	-8.163***	
LS	-0.970	-1.891	-5.258***	-5.219***	-0.948	-2.143	-5.277***	-5.242***	
MC	2.167	-1.957	-2.153	-3.707***	1.158	-1.005	-2.819**	-3.032***	
RICE	-1.974	-2.734	-7.479***	-7.465***	-1.974	-2.693	-8.129***	-8.141***	
СР	0.296	-2.051	-4.030***	-3.983**	-0.391	-2.802	-5.460	-5.365***	
ОТ	-2.161	-3.249	-7.307***	-0.614	-3.042	-2195	-6.787***	-6.789***	

Table 4a. Result of Ng-Perron test (At level).

Variable	Ng-Perron Test										
	At level										
	Constant				Constant with tr	end					
	MZa	MZt	MSB	MPT	MZa	MZt	MSB	MPT			
CO_2	-2.690	-0.989	0.368*	8.475*	-23.990	-3.448	0.144	3.891			
BM	-1.586	-0.645	0.407*	11.33*	-8.468	-2.044	0.241	10.803			
AG	1.449	1.338	0.923	65.343	-14.028	-2.648	0.189	6.500			
SF	0.119	0.089	0.749	35.161	-3.590	-1.213	0.338	23.335			
LS	1.026	0.781	0.761	43.220	-7.271	-1.893	0.260	12.554			
MC	-2.464	-0.793	0.322	8.401	-5.663	-1.611	0.284	15.923			
RICE	-2.782	-0.844	0.303	7.820	-13.354	-2.536	0.190	7.089			
CP	0.813	0.612	0.752	40.781	-15.360**	-2.763**	0.180	5.979			
OT	-6.755***	-1.773***	0.262	3.847	-8.515	-2.057	0.242*	10.721*			

Table 4b. Result of Ng-Perron test (At first difference).

	Ng-Perron Test							
Variable	At 1 st difference							
	Constant	Constant			Constant with tre			
	MZa	MZt	MSB	MPT	MZa	MZt	MSB	MPT
D(CO ₂)	-21.216*	-3.253*	0.153	1.169	-22.059**	-3.321**	0.151*	4.132*
D(BM)	-16.765*	-2.869*	0.171	1.557	-19.724**	-3.115**	0.158*	4.775*
D(AG)	-24.433*	-3.49*	0.143	1.021	-29.142**	-3.806**	0.131	3.190
D(SF)	-14.089*	-2.653*	0.188	1.743	-25.365*	-3.541*	0.140	3.714
D(LS)	-10.966**	-2.325**	0.212	2.299	-11.287	-2.374	0.210*	8.083*
D(MC)	-10.418**	-2.279**	0.219	2.364	-12.027	-2.449	0.204*	7.592*
D(RICE)	-6.951***	-1.82***	0.262	3.679	-13.183	-2.443	0.185*	7.587*
D(CP)	-23.729*	-3.410*	0.144	1.147	-23.648**	-3.423*	0.145*	3.944*
D(OT)	-23.629*	-3.432*	0.145	1.056	-25.338*	-3.558*	0.140	3.606

For Ng-Perron test, lag length is one. *, **, *** denotes the 1%, 5% and 10% level of significance, respectively.

Table 5. Cointegration result of Bayer-Hanck cointegration

Test statistic	EO-JOH	EO-JOH-BO-BDM	Decision
F-statistic	58.413	168.934	Cointegration Analysis
Critical regions			
1%	15	19.899	
5%	10.181	19.447	
10%	8.134	15.595	

F-statistic with consistent and decisive results. However, strong prohibition is needed of this hypothesis-based cointegration i.e. the order of each variable is stationary at first difference. An F-value greater than a critical value indicates that the null hypothesis of no cointegration should be rejected and F-values less than the critical value accepts the null hypothesis of no cointegration. Value of F-statistic under 1%, 5%, and 10% level of significance, both statistic is higher than critical values. Therefore, there exists a cointegration among variables.

4.4. Long run coefficients

As the variables were integrated at first order, the cointegration in this study was examined through FMOLS, DOLS, and CCR. Carbon dioxide emission is negatively associated with biomass-burned crop residues, agricultural land added to GDP, livestock, rice harvested area, cereal production, and other production in three models. From the data in Table 6, it can be seen that in FMOLS, the negative effect of CO₂ emissions is

significantly related to crop residues and biomass burning, livestock, rice harvested on land area, cereal production, and other crop production. This is because the more agricultural production there is, the more is the demand for use of combustible energy resources which consequently leads to release of emissions into the environment. This finding is similar to that of the studies conducted prior [86] and [87] in four ASEAN countries and in another study conducted [88] for E7 countries. CO2 emissions are pushed up a lot by the CO₂ equivalent of nitrous oxide (N₂O) emissions from synthetic fertilizers and other farming equipment [89]. In the case of DOLS, the results indicated that livestock and other crop production decreased CO₂ emissions. All variables' effects were statistically associated with CO2 emissions except crop residues and biomass burning in CCR. A study by Adedoyin et al. [88] found that energy use in the form of renewable energy had a negative and significant coefficient at various levels of significance. In the E7 countries, a 1% increase in energy use resulted in a 0.32 to 0.66 percent reduction in CO₂ emissions [88].

A causal connection was observed in Bayer-Hanck's cointegration. The Granger Causality test [90] was used to examine pairwise relationships. The results are presented in Table 7. Significant impact was observed on the agricultural value added to GDP when it comes to burning of biomass and crop residues and cereal production as well as environmental degradation and agricultural machinery. There was also a significant impact on the rice harvested area. Some unidirectional causality was observed from crop production to crop residues and biomass burning, livestock, rice harvested area, and other production (CP \rightarrow LS, CP \rightarrow RI, CP \rightarrow BM, CP \rightarrow OT). The results obtained in this study suggested that crop production was significantly affected by agricultural machinery, livestock plowed, rice cultivation, and other cereal production.

Table 6. Cointegrated coefficients of variables under FMOLS, DOLS and CCR.

Variable	FMOLS	FMOLS					CCR	CCR	
	Coefficient	t-statistic	p-value	Coefficient	t-Statistic	p-value	Coefficient	t-Statistic	p-value
BM	-2.295	-2.813	0.010	-6.017	-1.181	0.249	-0.866	-0.868	0.394
AG	-0.009	-0.034	0.973	1.314	0.801	0.431	-1.416	-4.208	0.000
SF	2.195	12.974	0.000	3.238	3.354	0.003	1.293	4.725	0.000
LS	-23.577	-24.683	0.000	-23.870	-4.037	0.001	-15.000	-14.552	0.000
MC	39.786	25.627	0.000	41.840	4.441	0.000	27.863	14.727	0.000
RICE	-2.444	-5.274	0.000	-1.405	-0.498	0.623	-1.666	-3.768	0.001
CP	-1.539	-4.061	0.001	-1.302	-0.566	0.576	-1.825	-2.877	0.009
OT	-0.630	-6.699	0.000	-1.365	-2.325	0.029	-0.481	-3.718	0.001
Constant	146.910	11.000	0.000	167.677	2.015	0.055	87.596	6.209	0.000

Table 7. Granger causality test.

Null Hypothesis	F- Statistic	p- value	Null Hypothesis	F- Statistic	p- value
AG does not Granger Cause CO ₂	8.708	0.001	AG does not Granger Cause RI	11.107	0.000
LS does not Granger Cause CO ₂	5.370	0.011	AG does not Granger Cause CP	12.374	0.000
CO ₂ does not Granger Cause LS	6.719	0.004	AG does not Granger Cause OT	4.269	0.025
MC does not Granger Cause CO ₂	12.702	0.000	SF does not Granger Cause RI	4.792	0.017
RI does not Granger Cause CO ₂	4.084	0.029	MC does not Granger Cause LS	6.483	0.005
CO ₂ does not Granger Cause RI	3.934	0.032	LS does not Granger Cause RI	4.616	0.019
CP does not Granger Cause CO ₂	8.436	0.002	CP does not Granger Cause LS	7.651	0.002
AG does not Granger Cause BM	7.048	0.004	RI does not Granger Cause MC	3.444	0.047
LS does not Granger Cause BM	3.436	0.047	MC does not Granger Cause RI	5.346	0.011
MC does not Granger Cause BM	3.526	0.044	MC does not Granger Cause OT	7.800	0.002
CP does not Granger Cause BM	4.604	0.019	CP does not Granger Cause RI	7.548	0.003
LS does not Granger Cause AG	7.774	0.002	RI does not Granger Cause OT	4.314	0.024
AG does not Granger Cause MC	7.203	0.003	CP does not Granger Cause OT	3.460	0.047
Null Hypothesis	F- Statistic	p- value	Null Hypothesis	F- Statistic	p- value

Unidirectional causality was observed for $CP \rightarrow CO_2$, $AG \rightarrow CO_2$, $MC \rightarrow CO_2$, $CP \rightarrow CO_2$ & $CP \rightarrow CO_2$, $LS \rightarrow BM$, $MC \rightarrow BM$, $AG \rightarrow BM$, $SF \rightarrow RI$, $AG \rightarrow RI$, $CP \rightarrow RI$, $MC \rightarrow RI$, and so on.

 CO_2 emissions and/or environmental degradation caused a 5% level of significance with livestock and rice harvested area in a bidirectional causality ($CO_2 \leftrightarrow LS$, $CO_2 \leftrightarrow RI$), which was similar to the findings of an earlier study [91], and suggests that as livestock increases, the chance of environmental degradation also increases. Rice is one of the staple foodgrains and the area where rice paddy fields are cultivated have a direct link to carbon emissions. Meijide et al. [92] claim that rice paddy fields are fundamental indicators of anthropogenic CO_2 emissions. Environmental degradation forces causal linkage of two factors into unidirectional causalities, such as agricultural machinery and cereal production ($MC \rightarrow CO_2$ & $CP \rightarrow CO_2$). Similar findings were also observed by Ullah et al. [60]. Therefore, it can be stated based on the observed findings that tillage and equipment used in an excessive manner can potentially stimulate environmental degradation. In Canadian firm, agricultural machinery and fuel play a role in CO_2 emissions [93]. Unlike the findings of [94] and [95], there

was no causal evidence of crop residue and biomass burning with CO_2 emissions increase. Crop production has been shown to cause CO_2 emissions increase in a unidirectional causality for Ghana [39]. Therefore, crop production can also positively attribute to carbon emissions. The monotonous increasing trend of environmental degradation can hamper the ecological balance. This onset triggers climatic hazards resulting from floods, cyclones, temperature increases, precipitation imbalances, and production in cereal areas being disrupted [96]. The findings of that study was opposite to that of Pant [97] who proposed that rising CO_2 can be utilized for agricultural productivity via a combination of accelerated photosynthesis and water use efficiency.

Several previous studies have studied the nexus between agriculture value addition, energy use, economic growth, and emissions in the context of different countries [60, 88, 98, 99, 100, 101]. However, this study differs from the existing literature by complementing it and considering the relationship between agricultural ecosystems and Bangladesh's environmental degradation from 1985 to 2017. Consequently, this study would be a major step forward in environmental and agricultural research. The findings of this study establishes a bidirectional causal relationship between environmental degradation and livestock, as well as between environmental degradation and rice harvest areas. This finding was previously supported by Owusu and Asumadu-Sarkodie [91], who discovered bidirectional causality between milled rice production and carbon dioxide emissions in their study. The results obtained in this study showed that environmental degradation had a one-way causal relationship with agricultural machinery and with cereal production. Another study in Pakistan [48] found similar results, stating that burning crop residues, livestock, farm equipment, cereal production, and other crop products all contribute significantly to CO2 emissions. Finally, the findings of the current study indicated that livestock, cereal production, agricultural machinery use, and agricultural land production all contributed to carbon dioxide emissions. Due to high land fragmentation and population growth, Bangladesh's cultivable land area has decreased over time, putting enormous pressure on the country's limited land resources to produce enough food for sustaining the growing population.

5. Limitations of the current study and scope for future research

Though there are various sources of anthropogenic emissions, this study focused only on the emissions resulting from agricultural production and agro-economy of Bangladesh. To achieve the proposed UNSDGs by 2030, any country must pay attention to social, economic, and environmental aspects. This study may support a two dimensional idea for policymakers to prepare for any future strategies. Moreover, the findings obtained in this study may provide an opportunity for future research to compare the situation of Bangladesh with that of other similar nations in order to identify whether obstacles are unique to all nations or vary from nation to nation. One limitation of this study was the lack of availability of most recent data and hence considered the datasets available until 2017.

6. Conclusions and policy implications

This study was confined to analysing the agricultural ecosystem from 1985 to 2017. On the other hand, rice as a staple food is considered in this study as a harvested area. Researchers consider more variables like cereal and other production, biomass burned crop residues, and agricultural machinery for modern technology to tackle the agroecosystem. The present study applied two customary unit root tests to study variables and concluded that each variable is stationary at I (1). Cointegration was confirmed by the Bayer-Hanck cointegration at a 1% level of significance. In particular the analysis of variable in FMOLS, DOLS, and CCR evidenced that environmental degradation is accelerated with a 1% significance in long-run estimates. A pairwise Granger Causality test was computed to check out causality with the results concluding that carbon dioxide emissions were caused by cereal production, agricultural production, and agricultural land production. Therefore, farming in Bangladesh is still considered to be causing environmental degradation due to unplanned and stagnant crop intensity combined with a lack of modern facilities. Due to high land fragmentation and a growing population, the arable land area of Bangladesh is declining, thereby creating huge pressure on limited land resources to produce sufficient food for all. To meet the nation's food requirements, the agricultural sector initiates such a method that is harmful to nature as well as our survival on earth. On the other hand, environmental degradation is accelerated due to unnecessary or unplanned farming of agricultural land, tillage use, lack of technological innovations and poor water management, primitive-based farming, securing employment, and raising entrepreneurialism from foreign investors and banking sides. This research envisages to contribute to a strong economic growth and a high-quality environment, keeping in line with the UN-SDG Targets of 7, 9, 12, and 13 on clean energy access and climate change mitigation. The majority of Bangladeshi farmers are illiterate and unaware of environmental issues. As a result, they harm nature both purposefully and accidently. Furthermore, most of the Bangladeshi population is not sufficiently conscious of environmental degradation, and there is a lack of environmental concern at every societal level. However, pollution can be reduced by implementing appropriate policies, laws, initiatives, and effective implementations. The findings of this study reveal that environmental contamination has a strong causal association with Bangladesh's agricultural ecology. By reinforcing and adapting the existing policies and strategies, the Government of Bangladesh should introduce some synchronized policies that may help achieve a sustainable agro-economic system by reducing environmental pollution. The following are some recom mendations.

- Increasing subsidies and government spending from a fiscal policy standpoint should result in certain projects that match demand.
- Establishing watchdog bodies to oversee the overall agriculture activities and identify the farmers responsible for creating more natural degradation.
- The use of harmful organic and inorganic products such as pesticides and chemical fertilizers should be reduced by developing awareness among farmers for this purpose and promoting biological evolution through using organic fertilizer.
- Technological advancement should be used to make the necessary stakeholders aware of the benefits of sustainable agriculture and protecting the environment.
- Though the farmers of Bangladesh are poor, setting penalties for fixed level of emissions transgressions may decrease environmental degradation.
- For a more robust agrarian economy, policies must be more aggressive for bridging the demand gap created by disasters or calamities that have devastated the low-lying lands.

Declarations

Author contribution statement

Shanjida Chowdhury: Conceived and designed the experiments; Analyzed and interpreted the data; Wrote the paper.

Sunjida Khan; Md Fouad Hossain Sarker; Md Kabirul Islam: Performed the experiments; Wrote the paper.

Maruf Ahmed Tamal: Analyzed and interpreted the data; Wrote the paper.

Niaz Ahmed Khan: Conceived and designed the experiments; Wrote the paper.

Funding statement

This research did not receive any specific grant from funding agencies in the public, commercial, or not-for-profit sectors.

Data availability statement

Data will be made available on request.

Declaration of interest's statement

The authors declare no conflict of interest.

Additional information

No additional information is available for this paper.

References

- M. Balat, H. Balat, N. Acici, Environmental issues relating to greenhouse carbon dioxide emissions in the world, Energy Explor. Exploit. 21 (5) (2003) 457–473.
- [2] United Nations Conference on Environment and Development, Rio de Janeiro, Brazil, June 1992, pp. 3–14.
- [3] B. Gyamfi, I. Ozturk, M. Bein, F. Bekun, An investigation into the anthropogenic effect of biomass energy utilization and economic sustainability on environmental degradation in E7 economies, Biofuels, Bioproducts and Biorefining 15 (3) (2021) 840–851.
- [4] A. Ali, Technological change in agriculture and land degradation in Bangladesh: a case study, Land Degrad. Dev. 15 (3) (2004) 283–298.
- [5] The Future of Food and Agriculture Trends and Challenges, Rome, 2017. Avaiable: https://www.fao.org/3/i6583e/i6583e.pdf.
- [6] [A. Ali, Technological change in agriculture and land degradation in Bangladesh: a case study, Land Degrad. Dev. 15 (3) (2004) 283–298.
- [7] FAO, The Future of Food and Agriculture Trends and Challenges. Rome, 2017. Available: https://www.fao.org/3/i6583e/i6583e.pdf.
- [8] D. Cervantes-Godoy, J. Dewbre, Economic importance of agriculture for poverty reduction, OECD Food, Agriculture and Fisheries Working Papers 23 (2010).
- [9] E. Hodgson, Toxicology and Human Environments, first ed., Elsevier Science, 2012
- [10] P. Kopittke, N. Menzies, P. Wang, B. McKenna, E. Lombi, Soil and the intensification of agriculture for global food security, Environ. Int. 132 (2019) 105078.
- [11] N. Khan, Climate Change and Sustainability of Agriculture in Bangladesh, Springer Climate, 2021, pp. 65–84.
- [12] F. Nunes, L. de Jesus Alves, C. de Carvalho, E. Gross, T. de Marchi Soares, M. Prasad, Soil as a complex ecological system for meeting food and nutritional security, Climate Change and Soil Interactions (2020) 229–269.
- security, Climate Change and Soil Interactions (2020) 229–269.

 [13] W. Aktar, D. Sengupta, A. Chowdhury, Impact of pesticides use in agriculture: their benefits and hazards, Interdiscipl. Toxicol. 2 (1) (2009) 1–12.
- [14] Data.worldbank.org, Population Density (People Per Sq. Km of Land Area) | Data, 2020 [Online]. Available: https://data.worldbank.org/indicator/EN.POP.DNST? locations=BD [Accessed: 19- Apr- 2022].
- [15] Bangladesh Climate Change Data | Emissions and Policies", Climatewatchdata.org. [Online]. Available: https://www.climatewatchdata.org/countries/BGD?end_year=2018&source=CAIT&start_year=1990. [Accessed: 19- Apr- 2022].
- [16] S. Pagiola, Environmental and natural resource degradation in intensive agriculture in Bangladesh, World Bank published in collaboration with Agriculture and Natural Resource Operations Division South Asia Region, Washington, D.C., 1005
- [17] R. Kanianska, Agriculture and Its Impacteon Land-Use, Eevironment, and Ecosystim Services, LandscapepEcology - bhe Influencea of Land Use and Anthropogenic Impacts of Landscape Creation (2016).

- [18] G. Jiang, et al., Effect of Agricultural Employment and Export Diversification Index on Environmental Pollution: Building the Agenda towards Sustainability, Sustainability 14 (2) (2022) 677.
- [19] Z. Ferdous, F. Zulfiqar, A. Datta, A. Hasan, A. Sarker, Potential and challenges of organic agriculture in Bangladesh: a review, J. Crop Improv. 35 (3) (2020) 403–426.
- [20] P. Kwakwa, V. Acheampong, S. Aboagye, Does agricultural development affect environmental quality? The case of carbon dioxide emission in Ghana, Manag. Environ. Qual. Int. J. 33 (2) (2021) 527–548.
- [21] M. Alam, R. Bell, W. Biswas, Increases in soil sequestered carbon under conservation agriculture cropping decrease the estimated greenhouse gas emissions of wetland rice using life cycle assessment, J. Clean. Prod. 224 (2019) 72-87
- [22] Noor-E-Sabiha, S. Rahman, Environment-smart agriculture and mapping of interactions among environmental factors at the farm level: a directed graph approach, Sustainability 10 (5) (2018) 1580.
- [23] N. Sabiha, R. Salim, S. Rahman, M. Rola-Rubzen, Measuring environmental sustainability in agriculture: a composite environmental impact index approach, J. Environ. Manag. 166 (2016) 84–93.
- [24] M.A.A. Faroque, M.A. Kashem, S.E. Bilkis, Sustainable agriculture: a challenge in Bangladesh, Int. J. Agric. Res. Innovat. Technol. 1 (1-2) (2011) 1–8.
- [25] Independent Group of Scientists Appointed by the Secretary-General, Global Sustainable Development Report 2019: the Future Is Now – Science for Achieving Sustainable Development, United Nations, New York, 2019. Available: https://sustainabledevelopment.un.org/content/documents/24797GSDR_report_2019.pdf.
- [26] Independent Group of Scientists Appointed by the Secretary-General, Global Sustainable Development Report 2019: the Future Is Now – Science for Achieving Sustainable Development, United Nations, New York, 2019. Available: https://sustainabledevelopment.un.org/content/documents/24797GSDR_report_2019.pdf.
- [27] T. Md, R. F, A. M, K. M, S. L, Relationship between environment and agriculture: a review, SunText Review of BioTechnology 1 (2) (2020).
- [28] S. Nathaniel, N. Adeleye, Environmental preservation amidst carbon emissions, energy consumption, and urbanization in selected african countries: implication for sustainability, J. Clean. Prod. 285 (2021) 125409.
- [29] M. Usman, S. Anwar, M. Yaseen, M. Makhdum, R. Kousar, A. Jahanger, Unveiling the dynamic relationship between agriculture value addition, energy utilization, tourism and environmental degradation in South Asia, J. Publ. Aff. (2021).
- [30] M. Hafeez, C. Yuan, W. Shah, M. Mahmood, X. Li, K. Iqbal, Evaluating the relationship among agriculture, energy demand, finance and environmental degradation in one belt and one road economies, Carbon Manag. 11 (2) (2020) 139–154.
- [31] I. Olanipekun, G. Olasehinde-Williams, R. Alao, Agriculture and environmental degradation in Africa: the role of income, Sci. Total Environ. 692 (2019) 60–67.
- [32] Danish M. Baloch, N. Mahmood, J. Zhang, Effect of natural resources, renewable energy and economic development on CO₂ emissions in BRICS countries, Sci. Total Environ. 678 (2019) 632–638.
- [33] T. Ramachandra, B. Aithal, K. Sreejith, GHG footprint of major cities in India, Renew. Sustain. Energy Rev. 44 (2015) 473–495.
- [34] D. Li, D. Yang, Does non-fossil energy usage lower CO₂ emissions? Empirical evidence from China, Sustainability 8 (9) (2016) 874.
- [35] H. Gholipour Fereidouni, Foreign direct investments in real estate sector and CO₂ emission, Manag. Environ. Qual. Int. J. 24 (4) (2013) 463–476.
- [36] A. Gattinger, J. Jawtusch, A. Muller, Mitigating Greenhouse Gases in Agriculture: A challenge and Opportunity for Agricultural Policies, Diakonisches Werk der EKD e. V. 2011
- [37] W. Schmidt, Powering the people beyond 2050, Sci. Technol. Dev. 31 (2) (2012) 133–151.
- [38] X. Liu, S. Zhang, J. Bae, The nexus of renewable energy-agriculture-environment in BRICS, Appl. Energy 204 (2017) 489–496.
- [39] S. Asumadu-Sarkodie, P. Owusu, Recent evidence of the relationship between carbon dioxide emissions, energy use, GDP, and population in Ghana: a linear regression approach, Energy Sources B Energy Econ. Plann. 12 (6) (2017) 495–503.
- [40] S. Ali, et al., Climate change and its impact on the yield of major food crops: evidence from Pakistan, Foods 6 (6) (2017) 39.
- [41] J. Pretty, Can sustainable agriculture feed Africa? New evidence on progress, processes and impacts, Environ. Dev. Sustain. 1 (34) (1999) 253–274.
- [42] G. Conway, The Doubly Green Revolution, Penguin Books, 1997.
- [43] J. Roberts, P. Grimes, J. Manale, Social roots of global environmental change: a world-systems analysis of carbon dioxide emissions, J. World Syst. Res. (2003) 277–315
- [44] F. Mariantonietta, S. Alessia, C. Francesco, P. Giustina, GHG and cattle farming: CO-assessing the emissions and economic performances in Italy, J. Clean. Prod. 172 (2018) 3704–3712.
- [45] S. Vetter, et al., Greenhouse gas emissions from agricultural food production to supply Indian diets: implications for climate change mitigation, Agric. Ecosyst. Environ. 237 (2017) 234–241.
- [46] J. Pretty, Agricultural sustainability: concepts, principles and evidence, Phil. Trans. Biol. Sci. 363 (1491) (2007) 447–465.
- [47] T. Aung, B. Saboori, E. Rasoulinezhad, Economic growth and environmental pollution in Myanmar: an analysis of environmental Kuznets curve, Environ. Sci. Pollut. Control Ser. 24 (25) (2017) 20487–20501.
- [48] D. Nezahat, Agriculture and Environmental Kuznets Curves in the case of Turkey: evidence from the ARDL and bounds test, Agric. Econ. 62 (12) (2016) 566–574.
- [49] Bangladesh | FAO regional office for asia and the pacific | food and agriculture organization of the united nations [Online]. Available: Fao.org (2022) [Accessed:

- $19-\ Apr-\ 2022], \ https://www.fao.org/asiapacific/perspectives/agricultural-statistics/global-strategy/results-in-the-region/bangladesh/en/.$
- [50] Bangladesh GDP From Agricultural ", Tradingeconomics.com. [Online]. Available: https://tradingeconomics.com/bangladesh/gdp-from-agriculture. [Accessed: 19-Apr- 2022].
- [51] Climate transparency report Bangladesh's climate action and responses to the COVID-19 crisis, Clim. Transp. (2020). Retrieved 18 April 2022, from, https:// www.climate-transparency.org/.
- [52] M. Hossain, S. Chen, The decoupling study of agricultural energy-driven CO₂ emissions from agricultural sector development, Int. J. Environ. Sci. Technol. (2021).
- [53] S. Rahman, Environmental consequences of agricultural development in Bangladesh: empirical evidence, farmer's perceptions and their determinants, in: J. Meijer, A. der Berg (Eds.), Handbook of Environmental Policy, Nova Science Publishers, New York, 2008.
- [54] W. Murad, J. Ratnatunga, Carbonomics of the Bangladesh agricultural output: causality and long-run equilibrium, Manag. Environ. Qual. Int. J. 24 (2) (2013) 256–271 [Accessed 19 April 2022].
- [55] B. Gunter, The impact of development on CO₂ emissions: a case study for Bangladesh until 2050, SSRN Electron. J. (2010).
- [56] Food and Agriculture Organization of the United Nations (FAO), 2017. Avaiable, http://www.fao.org/faostat/en/-data/QC.
- [57] S. Bhuvaneshwari, H. Hettiarachchi, J. Meegoda, Crop residue burning in India: policy challenges and potential solutions, Int. J. Environ. Res. Publ. Health 16 (5) (2019) 832.
- [58] I. Gurbuz, E. Nesirov, G. Ozkan, Does agricultural value-added induce environmental degradation? Evidence from Azerbaijan, Environ. Sci. Pollut. Control Ser. 28 (18) (2021) 23099–23112. Available:.
- [59] B. Hungate, et al., Assessing the effect of elevated carbon dioxide on soil carbon: a comparison of four meta-analyses, Global Change Biol. 15 (8) (2009) 2020–2034.
- [60] A. Ullah, D. Khan, I. Khan, S. Zheng, Does agricultural ecosystem cause environmental pollution in Pakistan? Promise and menace, Environ. Sci. Pollut. Control Ser. 25 (14) (2018) 13938–13955.
- [61] World development report 2017, Infrastructure for Development, Oxford University Press, New York, 2017.
- [62] C. Kao and M. Chiang, "On the estimation and inference of a cointegrated regression in panel data", Advances in Econometrics, pp. 179–222. DOI: https://doi.org/10.1016/S0731-9053(00)15007-8.
- [63] P. Pedroni, "Fully modified OLS for heterogeneous cointegrated panels", Adv. Econom., pp. 93-130. DOI: https://doi.org/10.1016/S0731-9053(00)15004-2.
- [64] A. Bashier, A. Siam, Immigration and economic growth in Jordan: FMOLS approach, International Journal of Humanities Social Sciences and Education 1 (9) (2014) 85–92 [Accessed 19 April 2022].
- [65] S. Narayan, P. Narayan, Determinants of demand for Fiji'S exports: an empirical investigation, Develop. Econ. 42 (1) (2004) 95–112.
- [66] D. Dickey, W. Fuller, Likelihood ratio statistics for autoregressive time series with a unit root, Econometrica 49 (4) (1981) 1057.
- [67] P. Phillips, P. Perron, Testing for a unit root in time series regression, Biometrika 75 (2) (1988) 335–346.
- [68] S. Johansen, K. Juselius, Maximum likelihood estimation and inference on cointegration - with applications to the demand for money, Oxf. Bull. Econ. Stat. 52 (2) (2009) 169–210.
- [69] R. Engle, C. Granger, Co-integration and error correction: representation, estimation, and testing, Econometrica 55 (2) (1987) 251. Available:.
- [70] H. Boswijk, Testing Stability and Identifiability of Long-Run Equilibria, 1994. Tinbergen Institute discussion paper, TI 94-101. [online] Amsterdam: Tinbergen Institute. Available at: https://hdl.handle.net/11245/1.423641 [Accessed 20 April 2022].
- [71] A. Banerjee, J. Dolado, R. Mestre, Error-correction mechanism tests for cointegration in a single-equation framework, J. Time Anal. 19 (3) (1998) 267–283. Available:.
- [72] E. Pesavento, Analytical evaluation of the power of tests for the absence of cointegration, J. Econom. 122 (2) (2004) 349–384.
- [73] C. Bayer, C. Hanck, Combining non-cointegration tests, J. Time Anal. 34 (1) (2013) 83–95.
- [74] Bruce E. Hansen, Peter C.B. Phillips, Estimation and Inference in Models of Cointegration: A Simulation Study, 1988. Cowles Foundation Discussion Papers. 1125, https://elischolar.library.yale.edu/cowles-discussion-paper-series/1125. Available:.
- [75] J. Stock, M. Watson, A simple estimator of cointegrating vectors in higher order integrated systems, Econometrica 61 (4) (1993) 783.
- 76] P. Adom, K. Amakye, C. Barnor, G. Quartey, The long-run impact of idiosyncratic and common shocks on industry output in Ghana, OPEC Energy Review 39 (1) (2015) 17–52.
- [77] Z. Rahman, M.E. Hossain, Role of agriculture in economic growth of Bangladesh: a VAR approach, Journal of Business Studies 7 (2014) 163–185.
- [78] F. Agbola, Does human capital constrain the impact of foreign direct investment and remittances on economic growth in Ghana? Appl. Econ. 45 (19) (2013) 2853–2862.
- [79] P. Adom, Asymmetric impacts of the determinants of energy intensity in Nigeria, Energy Econ. 49 (2015) 570–580.
- [80] J. Park, Canonical cointegrating regressions, Econometrica 60 (1) (1992) 119.
- [81] S. Naseem, T. Guang Ji, U. Kashif, M. Arshad, Causal analysis of the dynamic link between energy growth and environmental quality for agriculture sector: a piece of evidence from India, Environ. Dev. Sustain. 23 (5) (2020) 7913–7930.

- [82] A. Ghimire, F. Lin, P. Zhuang, The impacts of agricultural trade on economic growth and environmental pollution: evidence from Bangladesh using ARDL in the presence of structural breaks, Sustainability 13 (15) (2021) 8336.
- [83] D. DeJong, J. Nankervis, N. Savin, C. Whiteman, Integration versus trend stationary in time series, Econometrica 60 (2) (1992) 423.
- [84] R. Harris, R. Sollis, Applied Time Series Modelling and Forecasting, Wiley, 2003.
- [85] M. Chaouachi, D. Balsalobre-Lorente, Environmental strategies for achieving a new foreign direct investment golden decade in Algeria, Environ. Sci. Pollut. Control Ser. (2022).
- [86] M. Ben Jebli, S. Ben Youssef, Renewable energy consumption and agriculture: evidence for cointegration and Granger causality for Tunisian economy, Int. J. Sustain. Dev. World Ecol. 24 (2) (2016) 149–158.
- [87] X. Liu, S. Zhang, J. Bae, The nexus of renewable energy-agriculture-environ\ment in BRICS, Appl. Energy 204 (2017) 489–496.
- [88] F. Adedoyin, M. Bein, B. Gyamfi, F. Bekun, Does agricultural development induce environmental pollution in E7? A myth or reality, Environ. Sci. Pollut. Control Ser. 28 (31) (2021) 41869–41880.
- [89] P. Ciais, et al., Empirical estimates of regional carbon budgets imply reduced global soil heterotrophic respiration, Natl. Sci. Rev. 8 (2) (2020).
- [90] C. Granger, Investigating causal relations by econometric models and crossspectral methods, Econometrica 37 (3) (1969) 424.
- [91] P. Owusu, S. Asumadu-Sarkodie, Is there a causal effect between agricultural production and carbon dioxide emissions in Ghana? Environmental Engineering Research 22 (1) (2016) 40–54.
- [92] A. Meijide, C. Gruening, I. Goded, G. Seufert, A. Cescatti, Water management reduces greenhouse gas emissions in a Mediterranean rice paddy field, Agric. Ecosyst. Environ. 238 (2017) 168–178.

- [93] J. Dyer, R. Desjardins, Simulated farm fieldwork, energy consumption and related greenhouse gas emissions in Canada, Biosyst. Eng. 85 (4) (2003) 503–513. Available:.
- [94] A. Awasthi, S. Chauhan, S. Goyal, A fuzzy multicriteria approach for evaluating environmental performance of suppliers, Int. J. Prod. Econ. 126 (2) (2010) 370–378. Available:
- [95] B. Stanemann, et al., Mitigating the Impact of Agriculture on Air Quality and Climate Change: Solutions for Improved Nitrogen Management", IFOAM EU Group, B-Brussels, 2014.
- [96] A. Kumar, A. Nayak, S. Mohanty, B. Das, Greenhouse gas emission from direct seeded paddy fields under different soil water potentials in Eastern India, Agric. Ecosyst. Environ. 228 (2016) 111–123.
- [97] K. Pant, Effects of agriculture on climate change: a cross country study of factors affecting carbon emissions, Journal of Agriculture and Environment 10 (2009) 84–102.
- [98] K. Latif, M. Raza, Shahid adil and rehana kouser, "nexus between economy, agriculture, population, renewable energy and CO₂ emissions: evidence from asiapacific countries, Journal of Business and Social Review in Emerging Economies 6 (1) (2020) 261–276.
- [99] H. Mahmood, T. Alkhateeb, M. Al-Qahtani, Z. Allam, N. Ahmad, M. Furqan, Agriculture development and ${\rm CO_2}$ emissions nexus in Saudi Arabia, PLoS One 14 (12) (2019) e0225865.
- [100] S. Ghosh, K. Kanjilal, Long-term equilibrium relationship between urbanization, energy consumption and economic activity: empirical evidence from India, Energy 66 (2014) 324–331.
- [101] W. Xuezhen, W. Shilei, G. Feng, The Relationship between Economic Growth and Agricultural Growth: the Case of China, International Conference on E-Business and E-Government, 2010, pp. 5315–5318.